Mar Vista Animal Medical Center

3850 Grand View Blvd., Los Angeles, CA 90066

(310) 391-6741

Fax: (310) 391-6744

AMOXICILLIN

STRENGTH: 50 mg and 500 mg Capsules, 40 mg, 50 mg,100 mg, 125 mg, 150 mg, 200 mg, 250 mg, 400, 500, 875 mg, and Oral Suspension

ROUTE OF ADMINISTRATION: ORAL

May be given with or without food.

ACTION OF MEDICATION: Antibiotic. Best activity against Gram positive bacteria except Staphylococci.

MOST COMMON SIDE EFFECTS REPORTED: Mild upset stomach (vomiting, diarrhea or appetite loss). This is generally mitigated by giving the medication with food.

SEVERE SIDE EFFECTS REPORTED: The FDA has received sporadic reports of more serious reactions: fever, rashes, and even bone marrow suppression. These are extremely unusual but potentially very serious should they occur.

STORAGE: Keep away from light and store at room temperature. Oral suspension is best refrigerated.

IF YOU MISS A DOSE: Do not double up on the next dose. Simply give the medication when it is remembered or pick up with the next dose, allowing at least the proper interval between doses according to the label instructions.

OTHER INFORMATION: Other antibiotics can be enhanced by concurrent use with amoxicillin.

For a more detailed presentation, visit:

https://www.marvistavet.com/amoxicillin.pml