



Mar Vista Animal Medical Center

3850 Grand View Blvd., Los Angeles, CA 90066 (310) 391-6741 Fax: (310) 391-6744

COLCHICINE **(Brand Names: Colcrys, also commonly compounded individually)**

STRENGTHS: 0.6mg Capsules

ROUTE OF ADMINISTRATION: ORAL,
CAN BE GIVEN WITH OR WITHOUT FOOD,
SEE LABEL FOR INSTRUCTIONS.

USE ONLY IN DOGS!

ACTION OF MEDICATION: Inhibits collagenase to break down and prevent scarring, inhibits cell division.

MOST COMMON SIDE EFFECTS REPORTED: Vomiting, diarrhea, appetite loss. If seen, these symptoms can indicate the need for dose revision. Discontinue the medication and contact your veterinarian.

SERIOUS SIDE EFFECTS THAT HAVE BEEN REPORTED: Bone marrow suppression, Kidney toxicity.

STORAGE: Keep away from light. Store at room temperature.

IF YOU MISS A DOSE: Do not double up on the next dose. If less than 8 hours have passed since the dose was due, administer as normal. If more than 8 hours have passed, pick up with the next dose.

OTHER INFORMATION:

Drug interactions are reported with Cyclosporine, Azole antifungals, Chloramphenicol, Erythromycin, Diltiazem, and Cancer Chemotherapy Agents.

For a more detailed presentation, visit:

www.marvistavet.com/colchicine.pml