CYCLOPHOSPHAMIDE

(BRAND NAME: Cytoxan)

STRENGTH: 25 mg and 50 mg Tablets

ROUTE OF ADMINISTRATION: ORAL.

Should be given with food.

ACTION OF MEDICATION: Nitrogen Mustard derivative. Alkylating agent. DNA poison, used in the treatment of cancer. Standard doses are given every 3-4 weeks, "metronomic" therapy is given in a microdose daily.

MOST COMMON SIDE EFFECTS REPORTED: Vomiting, Diarrhea, hair loss in some breeds. Upset stomach is usually manageable with medication.

SEVERE SIDE EFFECTS REPORTED: Bone marrow suppression (can manifest as bruising but asymptomatic serious infections can be present). Marrow suppression occurs 5-14 days after administration. Hemorrhagic cystitis (bloody urine)

STORAGE: Keep away from light and store at room temperature. Do not refrigerate. Wear gloves when handling.

IF YOU MISS A DOSE: Check with your veterinarian for instructions. Never double up on the next dose.

OTHER INFORMATION: For several days after administration of this product, any body fluids or waste produced by the animal should be collected and disposed of in a plastic bag along with the gloves used for administration and for waste collection. This bag may be disposed of in the regular trash.

Do not split the tablets.

Pregnant women should not be exposed to this product.

For a more detailed presentation, visit:

www.marvistavet.com/cytoxan.pml