



3850 Grand View Blvd., Los Angeles, CA 90066 • (310) 391-6741 • Fax: (310) 391-6744  
<http://www.marvistavet.com>

# MARBOFLOXACIN

**(BRAND NAME: Zeniquin)**

**STRENGTH: 25 mg, 50 mg, 100 mg, 200 mg Tablets**

**ROUTE OF ADMINISTRATION:** Oral.  
Best given without food.

**ACTION OF MEDICATION:** Quinolone Class Antibiotic.

**MOST COMMON SIDE EFFECTS REPORTED:** Diarrhea, reduced appetite, nausea.  
These are generally mild and mitigated by giving with a small amount of food.

**SEVERE SIDE EFFECTS REPORTED:** Cartilage damage can occur when administered to young growing animals. Seizures are possible. Can form urinary crystals in higher doses. Other quinolones can lead to retinal damage in cats. Marbofloxacin is considered lower risk for this than other quinolones but it is still possible so doses are generally kept low in cats.

**STORAGE:** Keep away from light and store at room temperature.

**IF YOU MISS A DOSE:** Do not double up on the next dose. Simply give the medication when it is remembered or pick up with the next dose, allowing at least the proper interval between doses according to the label instructions.

**OTHER INFORMATION:** Calcium and other minerals will interfere with absorption. Do not use cheese as a treat to help administer this medication.

Several drug interactions are possible. See link below for details.

For a more detailed presentation, visit:

[www.marvistavet.com/marbofloxacin.pml](http://www.marvistavet.com/marbofloxacin.pml)